

4-1BBL/TNFSF9, Human, Recombinant

货号 : PCK282

产品信息

别名	CD137L, TNLG5A, TNFSF9
物种	Human
表达宿主	E.coli
序列信息	MREGPELSPDDPAGLLDLRQGMFAQLVAQNVLLIDGPLSWYSDPGLAGVS LTGGLSYKEDTKELVVAKAGVYYVFFQLELRRVVAGEGSGSVSLALHLQP LRSAAGAAALALTVDLPPASSEARNSAFGFQGRLLHLSAGQRLGVHLHTEA RARHAWQLTQGATVLGLFRVTPEIPAGLPSRSE with polyhistidine tag at the C-terminus.
检索号	P41273.1
分子量	20.38 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce IL-8 secretion in human PBMCs. The ED50 for this effect is <0.7 ng/mL.

产品特性

纯度	>95% as determined by SDS-PAGE. Ni-NTA chromatography.
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

4-1BBL is a transmembrane cytokine that is part of the tumor necrosis factor (TNF) ligand family. Recombinant human 4-1BB ligand is intended for use in cell culture applications. 4-1BBL and its interaction with 4-1BB is involved in the antigen presenting process, proliferation of CD4 and CD8 positive T-cells, as well as cytokine secretion from T-cells.

SDS-PAGE

