

## IL-15/IL-T, Mouse, Recombinant

货号 : PCK305

### 产品信息

别名	IL-T
物种	Mouse
表达宿主	E.coli
序列信息	NWIDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSCCKVTAMNCFLELQVILHE YSNMTLNETVRNVLYLANSTLSSKNVAESGCKECELEEKTFTEFLQSFIR IVQMFINTS with polyhistidine tag at the N-terminus.
检索号	P48346.1
分子量	14.2 kDa
标签	His-tag at the N-terminus
生物活性	Measure by its ability to induce CTLL-2 cells proliferation. The ED50 for this effect is <10 ng/mL. The specific activity of recombinant mouse IL-15 is approximately >1x 10 <sup>5</sup> IU/mg.

### 产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



## 复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## 背景介绍

Interleukin-15 (IL-15) is a cytokine with structural similarity to Interleukin-2 (IL-2). Like IL-2, IL-15 binds to and signals through a complex composed of IL-2/IL-15 receptor beta chain (CD122) and the common gamma chain (gamma-C, CD132). IL-15 is secreted by mononuclear phagocytes (and some other cells) following infection by virus(es). This cytokine induces cell proliferation of natural killer cells; cells of the innate immune system whose principal role is to kill virally infected cells.

## SDS-PAGE

