

G-CSF/CSF-3/MGI-1G/pluripoietin, Mouse, Recombinant

货号 : PCK333

产品信息

别名	CSF-3, MGI-1G, GM-CSF beta, pluripoietin
物种	Mouse
表达宿主	E.coli
序列信息	VPLVTVSALPPSLPLPRSFLKSLQVRKIQASGSVLLEQLCATYKLCHPEEL VLLGHSLGIPKASLSGCSSQALQQTQCLSQLHSGLCCLYQGLLQALSGISPAL APTLDLLQLDVANFATTIWQQMENLGVAPTQPTQSAMPAFTSAFQRRAG GVLAISYLQGFLETARLALHHLA with polyhistidine tag at the N- terminus.
检索号	P09920.1
分子量	19.76 kDa
标签	His-tag at the N-terminus
生物活性	Measure by its ability to induce proliferation in NFS-60 cells. The ED50 for this effect is <50 pg/mL. The specific activity of recombinant mouse G-CSF is > 2 x 10 ⁷ IU/mg.

产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

G-CSF is a hematopoietic growth factor. It can activate the progress that committed progenitor cells develop to neutrophils and enhance the functional activities of the mature end-cell. It is secreted in response to specific stimulation by a variety of cells, including bone marrow stroma, macrophages, endothelial cells and fibroblasts. In clinical treatment, G-CSF is used to facilitate hematopoietic recovery after bone marrow transplantation.

SDS-PAGE

