

## CCL4/MIP-1 $\beta$ /ACT-2, Mouse, Recombinant

货号 : PCK336

### 产品信息

别名	MIP-1b: Macrophage Inflammatory Protein-1 $\beta$ , ACT-2
物种	Mouse
表达宿主	E.coli
序列信息	APMGSDPPTSCCFSYTSRQLHRSFVMDYYETSSLCSKPAVVFLTKRGRQICANPSEPWVTEYMSDLELN with polyhistidine tag at the N-terminus.
检索号	Q5QNV9
分子量	8.64 kDa
标签	His-tag at the N-terminus
生物活性	Measure by its ability to chemoattract BaF3 cells transfected with human CCR5. The ED50 for this effect is <4 ng/mL.

### 产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 $\mu$ g of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 $\mu$ m filtered solution containing 1X PBS, pH 7.4.



## 复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## 背景介绍

CCL4 act as an important pro-inflammatory cytokine during acute and chronic inflammatory responses. CCL4 also play a role as an attractant NK cells, dendritic cells, monocytes, and lymphocytes to sites of injury and inflammation. As CCL4 conducts signaling through G-protein-coupled receptor CCR5, which is a major co-receptor for M-tropic HIV strains. Thus, binding of CCL4 to CCR5 inhibits HIV entry and reduces the cell surface expression of CCR5. CCL4 has been identified as one of the major anti-HIV factors produced by CD8+ T cells.

## SDS-PAGE

