

TNF- β / TNFB/ TNFSF1, Human, Recombinant

货号 : PCK053

产品信息

别名	Lymphotoxin-Alpha; LT-Alpha; TNF-Beta; Tumor Necrosis Factor Ligand Superfamily Member 1; LTA; TNFB; TNFSF1
物种	Human
表达宿主	E.coli
序列信息	Leu35-Leu205
检索号	P01374
分子量	18.8 kDa
生物活性	Measured in a cytotoxicity assay using L-929 mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D. The ED50 for this effect is 20-80 pg/ml.

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per μ g as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

Tumor Necrosis Factor β (TNF- β) is a secreted Protein belonging to the tumor necrosis factor family. TNF- β binds to TNFRSF1A/ TNFR1, TNFRSF1B/ TNFBR and TNFRSF14/ HVEM in homotrimeric form, binds to TNFRSF3/ LTBR in heterotrimeric form with LTB. TNF- β forms heterotrimers with lymphotoxin-beta, which anchors TNF- β to the cell surface. TNF- β mediates the inflammatory, immunostimulatory, and antiviral response, involves in the formation of second lymphoid organs during development, has a role in apoptosis. TNF- β is produced by lymphocytes and cytotoxic for a variety of tumor cells in vitro and in vivo.

SDS-PAGE



生物活性



Measured in a cytotoxicity assay using L-929 mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D. The ED50 for this effect is 20-80 pg/ml.

