

IL-2RB/CD122 (C-6His), Human, Recombinant

货号:PCK151

产品信息

别名 Interleukin-2 Receptor subunit beta; IL2RB; IL-2 Receptor subunit beta; IL-2R

subunit beta; High affinity IL-2 Receptor subunit beta; CD122

物种 Human

表达宿主 Human Cells 🍸

序列信息 Ala27-Asp239

检索号 P14784

分子量 25.6 kDa

标签 C-6His

生物活性 Loaded Human IL-2RB-His on HIS1K Biosensor, can bind Human IL-2 with an

affinity constant of 0.27 uM as determined in BLI assay.

产品特性

纯度 >95% as determined by reducing SDS-PAGE.

内毒素 <1.0 EU per µg as determined by LAL test.

保存 Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at -5~-20°C for 3 months.

运输 Ambient temperature or ice pack.

制剂 Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

网站: <u>www.procell.com.cn</u> 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋





Rev. V1.0




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml.Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

Human IL-2RB, also known asInterleukin-2 Receptor subunit beta, is the Receptor for Interleukin-2. IL2 Receptor complex is involved in Receptor mediated endocytosis and transduces the mitogenic signals of IL2. IL2 Receptor complex has three forms with respect to ability to bind IL2. IL-2RB is belonged to a type I membrane Protein, and has a 26 residue signal peptide, a 214 residue extracellular region, a 25 residue transmembrane region and a 286 residue cytoplasmic domain. IL-2RB is the subunit critical for Receptor-mediated signaling via physically or functionally coupling to other signaling molecules, such as the Jak-STAT and Src-family Protein tyrosine kinase although it lacks apparent catalytic Motifs.

SDS-PAGE



生物活性



Loaded Human IL-2RB-His on HIS1K Biosensor, can bind Human IL-2 with an affinity constant of 0.27 uM as determined in BLI assay

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



