

# IGF-II/ IGF2, Human, Recombinant

货号: PCK155

### 产品信息

别名 Insulin-Like Growth Factor II; IGF-II; Somatomedin-A; IGF2; PP1446 الالمنية

物种 Human

表达宿主 E.coli

序列信息 Ala25-Glu91

检索号 P01344

分子量 8.9 kDa

# 产品特性

纯度 >95% as determined by reducing SDS-PAGE.

内毒素 <1.0 EU per µg as determined by LAL test.

保存 Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at -5~-20°C for 3 months.

运输 Ambient temperature or ice pack.

制剂 Lyophilized from a 0.2 µm filtered solution of 20mM Glycine-HCl, 4% Sucrose,

4% Mannitol, 0.02% Tween 80 (w/v), pH3.0.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋








#### 复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml.Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## 背景介绍

Insulin-Like Growth Factor II (IGF2) belongs to the Insulin family of polypeptide Growth Factors that is involved in development and growth. Members of this family are structurally homologous to proInsulin, and share higher sequence identity. IGF2 is expressed only from the paternally inherited allele and believed to be secreted by the liver and to circulate in the blood. IGF2 possess growthpromoting activity and can stimulate the proliferation and survival of various cell types including muscle, bone, and cartilage tissue in vitro. IGF2 is influenced by placental lactogen and may play a role in fetal development.





网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



