

IL-17A/CTLA-8, Human, Recombinant

货号 : PCK177

产品信息

别名	CTLA-8
物种	Human
表达宿主	E.coli
序列信息	MIVKAGITIPRNP GCPNSEDKNFPRTVMVNLNIHNRNTNTNPKRSSDYYNRS TSPWNLHRNEDPERYPSVIWEAKCRHLGCINADGNVDYHMNSVPIQQEILV LRREPPHCPNSFRLEKILVSVGCTCVTPIVHHVA with polyhistidine tag at the C-terminus.
检索号	Q16552.1
分子量	16.47 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED50 for this effect is <6 ng/mL.

产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.01 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g.0.1% BSA,10%FBS,5%HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 $\mu\text{g/mL}$. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin-17A is a protein that in humans is encoded by the IL17A gene. The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO).

SDS-PAGE

