

ANGPTL3 (Ser17-Thr206) (C-6His), Mouse, Recombinant

货号：PCK226

产品信息

别名	Angiopoietin-related Protein 3; Angiopoietin-like Protein 3; Angptl3
物种	Mouse
表达宿主	Human Cells
序列信息	Ser17-Thr206
检索号	Q9R182
分子量	22.7 kDa

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.



复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Angiopoietin-like Protein 3 (ANGPTL3) is a secreted glyco Protein that is structurally related to the angiopoietins. Mature mouse ANGPTL3 contains an N-terminal coiled coil domain and a C-terminal fibrinogen-like domain. Within the N-terminal fragment, mouse ANGPTL3 shares 83% and 92% aa sequence identity with human and rat ANGPTL3, respectively. ANGPTL3 is expressed in the liver from early in development through adulthood. ANGPTL3 directly inhibits lipoprotein lipase (LPL) and endothelial lipase (EL), enzymes responsible for hydrolyzing circulating triglycerides and HDL phospholipids. This activity requires a putative heparin-binding motif which is N-terminal to the coiled coil domain. Proteolytic removal of the fibrinogen-like domain from the N-terminal fragment serves to activate ANGPTL3 and increase its ability to inhibit LPL in vitro and function in vivo. ANGPTL3 promotes an increase in circulating triglyceride levels without altering VLDL or HDL secretion or uptake. ANGPTL3 expression in vivo is up-regulated by LXR agonists and down-regulated by insulin, leptin, and agonists of TRβ or PPARβ. ANGPTL3, secreted by fetal liver cells, also promotes the expansion of hematopoietic stem cells.

SDS-PAGE

